

# A Stereo-Atlas of Ostracod Shells

edited by R. H. Bate, J. W. Neale, Lesley M. Sheppard  
and David J. Siveter

Volume 7, Part 2; 19th December 1980

Published by The British Micropalaeontological Society  
in association with Robertson Research International Ltd., Llandudno, Wales

## Editors

- Dr. R. H. Bate, Department of Palaeontology, British Museum (Natural History), Cromwell Road, London SW7 5BD.  
Prof. J. W. Neale, Department of Geology, The University, Hull HU6 7RH.  
Ms. Lesley M. Sheppard, Department of Palaeontology, British Museum (Natural History), Cromwell Road, London SW7 5BD.  
Dr. David J. Siveter, Department of Geology, The University, Leicester LE1 7RH.

## Editorial Board

- Dr. Richard H. Benson, Smithsonian Institution, Washington, D.C., 20560. U.S.A.  
Dr. Alwine Bertels, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Argentina.  
Dr. K. Ishizaki, Institute of Geology and Paleontology, Tohoku University, Sendai, Japan.  
Dr. C. W. Haskins, Robertson Research International Limited, 'Ty'n-y-Coed', Llanrhos, Llandudno, N. Wales, LL30 1SA.  
Dr. P. J. Jones, Bureau of Mineral Resources, P.O. Box 378, Canberra City, A.C.T. 2601, Australia.  
Prof. Dr. E. Kempf, Geologisches Institut der Universität Köln, Zùlpicher Strasse 49, D-5 Köln 1, German Federal Republic.  
Dr. H. J. Oertli, S.N.P.A., Centre de Recherches, 64001 Pau, France.  
Prof. G. Ruggieri, Istituto e Museo di Geologia dell'Università di Palermo, Corso Tuköry, 131, 90134 Palermo, Italy.  
Mr. P. F. Sherrington, Petro-Canada, P.O. Box 2844, Calgary, Alberta T2P 2M7, Canada.

## Instructions to Authors

Contributions illustrated by scanning electron micrographs of Ostracoda in stereo-pairs are invited. Full instructions may be obtained on request from any one of the Editors or Editorial Board. Format should follow the style set by the majority of papers in this issue. Descriptive matter apart from illustrations should be cut to a minimum; preferably each plate should be accompanied by one page of text only. Blanks to aid in mounting figures for plates may be obtained from the Editors. Completed papers should be sent to Ms. L. M. Sheppard, Department of Palaeontology, British Museum (Natural History), Cromwell Road, London SW7 5BD.

## Acknowledgments

This Volume of the *Stereo-Atlas* has been aided by generous financial support from Robertson Research International Limited.

## Stereo-viewing for users of the Atlas

In order to obtain maximum information and benefit from the use of the *Stereo-Atlas* it is essential that the user view the micrographs stereoscopically. Small pocket-sized stereo-viewers are most suitable for this purpose. Two suppliers are:

C. F. Casella & Co. Ltd., Regent House, Britannia Walk, London N1 7ND and  
Air Photo Supply Corp., 158 South Station, Yonkers, New York 10705. U.S.A.

The front cover shows a ventral view of the Cretaceous *Pattersonocypris micropapilosa* Bate preserved with appendages.

Plates reproduced by Torr of Silverstone, Northants., England.



ON *BULBOSCLERITES UNICORNIS* (NECKAJA)

by Roger E. L. Schallreuter  
(University of Hamburg, German Federal Republic)

*Bulbosclerites unicornis* (Neckaja, 1952)

- 1952 *Mica unicornis* sp. nov. A. I. Neckaja, *Trudŭ vses neft nauchno-issled. geol. - razv. Inst.*, **60** (= *Mikrofauna SSSR* 5), 217, 229, 230, pl. 3, figs. 1, 2.  
1962 *Rectella unicornis* (Neckaja); L. I. Sarv, *Eesti NSV Tead. Akad. Geol. Inst. uurimused*, **9**, tab. 1.  
1967 *Rectella ? unicornis* (Neckaja); R. E. L. Schallreuter, *Geologie*, **16** (5), 616.  
1973 *Rectella unicornis* Neckaja; A. I. Neckaja, *Trudŭ vses neft nauchno-issled. geol. - razv. Inst.*, **324**, 70.  
1979 *Bulbosclerites unicornis* (Neckaja); R. E. L. Schallreuter, *Taxonomy, Biostratigraphy and Distribution of Ostracodes* (Ed. N. Krstiĉ), Proc. VII Internat. Symp. Ostracodes, 26.

*Holotype*: Vsesojuznyj neftjanoy nauĉno-issledovatel'skij geologorazvedoĉnyj institut (VNIGRI), Leningrad, no. 14 - 128, carapace.

*Type locality*: Raion of Kamariku, Estonia; approx. lat. 59° 10' N, long. 26° 10' E. Porkuni Stage (F<sub>2</sub>), uppermost Ordovician.

Explanation of Plate 7, 74

Fig. 1, LV, ext. lat. (GPIH 2343, 570 µm long); fig. 2, RV, ext. lat. (GPIH 2344, 620 µm long).

Scale A (100 µm; x 175), fig. 1; scale B (100 µm; x 155), fig. 2.

*Figured specimens*: Geologisch-Paläontologisches Institut, University of Hamburg (GPIH), nos. 2343 (LV: Pl. 7, 74, fig. 1), 2344 (RV: Pl. 7, 74, fig. 2), 2345 (car.: Pl. 7, 76, fig. 1) and 2346 (LV: Pl. 7, 76, fig. 2). All the figured specimens are from the Öjlemyrflint (Upper Ordovician) erratic boulder no. Sy 2 of the Kaolinsand (Pliocene-Pleistocene), near Braderup, Isle of Sylt (N Frisian Is., N Sea); lat. 54° 56' N, long. 8° 21' E; coll. by Ulrich von Hacht, 1976.

*Diagnosis*: Species of *Bulbosclerites* with length up to 0.70mm. Length/height ratio >2.00. Posterodorsal process of the right valve is projected as a spine.

*Remarks*: The type-species *B. longa* Knüpfer (*Freiberger ForschHft.*, ser. C, **234**, 1968) is larger (0.78mm) and relatively higher (length/height ratio <2.00) and its posterodorsal process on the right valve is more bulb-like rather than spine-like as in *B. unicornis*. The latter possesses stop pegs in the left valve (see Pl. 7, 76, fig. 2 and Schallreuter, 26, 1979).

The spine in the holotype of *B. unicornis* (Neckaja, *op. cit.*, pl. 3, figs. 1, 2) is considerably shorter than those in the specimens figured herein (Pl. 7, 74, fig. 2; Pl. 7, 76, fig. 1); it is possible that the material represents a distinct subspecies.

The assignment of *Parasclerites postnodus* Blumenstengel (*Ibid.*, **182**, 1965) to the genus is questionable because of the more ventral position of the relatively weak, broad, knob-like inflation of the right valve.


*Distribution*: Porkuni Stage (F<sub>2</sub>), uppermost Ordovician, of Estonia. Öjlemyrflint (Upper Ordovician) erratic boulders of the Isle of Gotland (Baltic Sea) and of the Kaolinsand (Pliocene - Pleistocene) of the Isle of Sylt (N Frisian Is., N Sea).

Explanation of Plate 7, 76

Fig. 1, car., ext. vent. obl. (GPIH 2345, 575 µm long); fig. 2, LV, int. dors. obl. (GPIH 2346, 590 µm long).

Scale A (100 µm; x 170), figs. 1, 2.











ON *PACHYDOMELLOIDES BRADERUPENSIS* SCHALLREUTER sp. nov.

by Roger E. L. Schallreuter  
(University of Hamburg, German Federal Republic)

*Pachydomelloides braderupensis* sp. nov.

**Holotype:** Geologisch-Paläontologisches Institut, University of Hamburg, no. 2460, LV.  
[Paratype: no. 2462, LV].

**Type locality:** Middle Ordovician Hornstein erratic boulder no. Sy 52 of the Kaolinsand (Pliocene – Pleistocene), near Braderup, Isle of Sylt (N Frisian Is, N Sea), Germany; lat. 54° 56' N, long. 8° 21' E.

**Derivation of name:** After the type locality, Braderup, Isle of Sylt.

**Figured specimens:** Geologisch-Paläontologisches Institut, University of Hamburg (GPIH) nos. 2459 (RV: Pl. 7, 78, fig. 1), 2460 (LV: Pl. 7, 78, fig. 2), 2461 (LV: Pl. 7, 80, fig. 1) and 2462 (LV: Pl. 7, 80, fig. 2). From the lower Upper Viruan (Middle Ordovician) Hornstein erratic boulders nos. Sy 39 (2461) and Sy 52 (2459, 2460, 2462) from the type locality; coll. by Ulrich von Hacht, 1978.

---

Explanation of Plate 7, 78

Fig. 1, RV, ext. lat. (GPIH 2459, 617 µm long); fig. 2, LV, ext. lat. (GPIH 2460, 636 µm long).  
Scale A (100 µm; x 145), fig. 1; scale B (100 µm; x 125), fig. 2.

**Diagnosis:** Species of *Pachydomelloides* with length up to 0.64mm (holotype). Posterior end relatively broadly rounded. Sulcament not observed, perhaps lacking.

**Remarks:** *P. valcourensis* Swain (*J. Paleo.*, 36 (4), 739, 1962), the type-species, and *P. imperfecta* (Krause) (*Z. Deutsch. geol. Ges.*, 48 (1896) (4), 935, 1897) are both much larger, being more than 1mm long. Both species possess an inner sulcament (subvertical ridge) which has not been observed in *P. braderupensis*, but this may be a factor of preservation or of infraspecific variation. Furthermore, in both species the posterior end of the left valve seems to be more acuminate (cf. Swain, *op. cit.*, pl. 111, fig. 8a) than in *P. braderupensis*. Stop pegs (stop ridges, internal prongs), present in *P. braderupensis* (Pl. 7, 80, figs. 1, 2), were not mentioned in the description of the type-species but were observed by Krause (*op. cit.*, 936, pl. 25, fig. 11) in *P. imperfecta*.



**Distribution:** Lower Upper Viruan (Middle Ordovician) Hornstein erratic boulders of the Kaolinsand (Pliocene – Pleistocene) near Braderup, Isle of Sylt (N Frisian Is, N Sea), Germany.

---

Explanation of Plate 7, 80

Fig. 1, LV, int. lat. (GPIH 2461, 622 µm long); fig. 2, LV, int. lat. (GPIH 2462, 633 µm long).  
Scale A (100 µm; x 135), fig. 1; scale B (100 µm; x 140), fig. 2.











ON *CYPRIDEIS UNDOSA* VAN HARTEN sp. nov.

by Dick van Harten  
(University of Amsterdam, The Netherlands)

*Cyprideis undosa* sp. nov.

1980 *Cyprideis* sp., L.P.A. Geerlings *et al.*, *Proc. K. ned. Akad. Wet.*, Amsterdam, ser. B, 83, 29 - 37 (*pars*).

*Holotype*: Geological Institute University of Amsterdam, coll. no. PA 8768, ♀ LV.

[Paratypes: Geological Institute, University of Amsterdam, coll. nos. PA 8769 - 8774].

*Type locality*: N bank of Rio Almanzora, approx. 5km NE of Vera, Province of Almería, Spain; Grid Ref.; UTM 300.5, 763.5 (see text-fig. 1).

Light greenish grey laminated marl with *Chara* and abundant brown reed-like plant remains; Messinian.

*Derivation of name*: Latin *undusus*, full of waves; alluding to the sinuous course of ventral margin.

---

Explanation of Plate 7, 82

Fig. 1, ♀ LV, ext. lat. (holotype, PA 8768, 890  $\mu$ m long); fig. 2, ♀ RV, ext. lat. (paratype, PA 8769, 880  $\mu$ m long); fig. 3, ♀ car., ext. dors. (paratype, PA 8770, 890  $\mu$ m long).

Scale A (250  $\mu$ m; x 70), figs. 1 - 3.

*Figured specimens*: Geological Institute, University of Amsterdam, coll. nos. PA 8768 (holotype, ♀ LV: Pl. 7, 82, fig. 1), PA 8769 (♀ RV: Pl. 7, 82, fig. 2), PA 8770 (♀ car.: Pl. 7, 82, fig. 3), PA 8771 (♀ LV: Pl. 7, 84, fig. 1), PA 8772 (♂ LV: Pl. 7, 86, fig. 1), PA 8773 (♂ RV: Pl. 7, 86, fig. 3), PA 8775 (♀ car.: Pl. 7, 84, fig. 2), PA 8776 (♀ RV: Pl. 7, 84, fig. 3; Pl. 7, 88, fig. 3), PA 8777 (♀ RV: Pl. 7, 86, fig. 2), PA 8778 (juv. - 1 RV: Pl. 7, 88, fig. 1), PA 8779 (juv. - 1 RV: Pl. 7, 88, fig. 2).

All figured material is from the type locality near Vera, Spain; Grid Ref. UTM 300.5, 763.5; Messinian, light greenish grey laminated marl with *Chara* and brown reed-like plant remains; coll. by H. Dronkert and L.P.A. Geerlings.

*Diagnosis*: Medium-sized species of *Cyprideis* with a sinuous ventral margin (notably so in the males) and conspicuous surface pitting. High nodding potential (see text-fig. 2). Anterior margins non-denticulate. Right valve may carry up to two posteroventral spines.

*Remarks*: As to size, ventral sinuosity and type of ornamentation, this species resembles *Cyprideis seminulum* (Reuss), in particular the subspecies *C. seminulum portaferricum* Krstić (see M. A. Bassiouni, *Geol. Jb. B* 31, 92, pl. 1, figs. 7 - 10, 1979). The two species significantly differ in the lateral outline of the posterior part of their shells, in their nodding characteristics and in the presence of anteromarginal denticulation. At the type locality *C. undosa* sp. nov. is sympatric with *C. agrigentina* Decima (*Paleont. Ital.* 57, 108, 1964) and *C. calabra* Decima (*op. cit.*, 127).

A shallow, oligohaline palaeoenvironment is inferred for the new species.



---

Explanation of Plate 7, 84

Fig. 1, ♀ LV, ext. lat. (paratype, PA 8771, 890  $\mu$ m long); fig. 2, ♀ car., ext. vent. (PA 8775, 930  $\mu$ m long); fig. 3, ♀ RV, ext. lat. (PA 8776, 790  $\mu$ m long).

Scale A (250  $\mu$ m; x 70), figs. 1, 2; scale B (250  $\mu$ m; x 71), fig. 3.












**Distribution:** Outside the type locality *C. undosa* has been found in beds of comparable age near Sorbas, SE Spain. It seems to be associated with the freshening conditions that succeeded the deposition of evaporites in

the Messinian of the western Mediterranean area (see T.B. Roep & D. van Harten, *Ann. Géol. Pays Hellén.*, Tome hors série, fasc. 3, 1037 - 1044, 1979 and Geerlings *et al.*, *op. cit.*).

The vertical range of the species is not known but probably rather restricted.




Text-fig. 1. Map of Vera area. Type locality indicated with arrow.

#### Explanation of Plate 7, 86

Fig. 1, ♂ LV, ext. lat. (paratype, PA 8772, 900  $\mu$ m long); fig. 2, ♀ RV, int. lat. (PA 8777, 860  $\mu$ m long); fig. 3, ♂ RV, ext. lat. (paratype, PA 8773, 930  $\mu$ m long).

Scale A (250  $\mu$ m; x 70), figs. 1, 2; scale B (250  $\mu$ m; x 65), fig. 3.



Text-fig. 2. Ext. lat. view of ♀ RV showing possible positions of nodes in *Cyprideis undosa* sp. nov. (A) as compared with Sandberg's general scheme (B). B after Sandberg (1964).



**Acknowledgement:** I am grateful to Hans Dronkert and Peter Geerlings who kindly provided the material on which this paper is based.

#### Explanation of Plate 7, 88

Fig. 1, juv. - 1 RV, int. lat. (PA 8778, 710  $\mu$ m long); fig. 2, juv. - 1 RV, ext. lat. (PA 8779, 710  $\mu$ m long); fig. 3, ♀ RV, ext. lat., central group of nodes (PA 8776).

Scale A (250  $\mu$ m, x 86), figs. 1, 2; scale B (100  $\mu$ m; x 172), fig. 3.











ON *CYPRIDEIS EXUBERANS* VAN HARTEN sp. nov.

by Dick van Harten  
(University of Amsterdam, The Netherlands)

*Cyprideis exuberans* sp. nov.

*Holotype*: Geological Institute, University of Amsterdam, coll. no. PA 8780, ♀ LV.  
[Paratypes: Geological Institute, University of Amsterdam, coll. nos. PA 8781 - 8786].

*Type locality*: N bank of Arroyo de Abejuela, W of bridge in road from Elche de la Sierra to Ferez, Province of Albacete, Spain; lat. 38° 22' N, long. 02° 03' W. Yellowish calcarenite with *Chara* and abundant small gastropods; lacustrine, Miocene — Pliocene.

*Derivation of name*: Latin *exuberans*, abounding; alluding to rather exaggerated nature of diagnostic features.

---

Explanation of Plate 7, 90

Fig. 1, ♀ LV, ext. lat. (holotype, PA 8780, 780 µm long); fig. 2, ♀ RV, ext. lat. (paratype, PA 8782, 730 µm long); fig. 3, ♀ car., ext. dors. (PA 8790, 790 µm long).  
Scale A (250 µm; x 75), figs. 1 - 3.

*Figured specimens*: Geological Institute, University of Amsterdam, coll. nos. PA 8780 (holotype, ♀ LV: Pl. 7, 90, fig. 1), PA 8782 (♀ RV: Pl. 7, 90, fig. 2), PA 8783 (♂ RV: Pl. 7, 92, fig. 2), PA 8784 (♂ LV: Pl. 7, 92, fig. 1), PA 8785 (♀ car.: Pl. 7, 94, fig. 3), PA 8786 (♂ car.: Pl. 7, 92, fig. 3), PA 8789 (♂ LV: Pl. 7, 96, fig. 4), PA 8790 (♀ car.: Pl. 7, 90, fig. 3), PA 8791 (juv. - 1 LV: Pl. 7, 98, fig. 2), PA 8792 (juv. - 1 LV: Pl. 7, 98, fig. 1), PA 8793 (juv. - 1 RV: Pl. 7, 100, fig. 2), PA 8794 (♀ LV: Pl. 7, 94, fig. 2), PA 8795 (♀ LV: Pl. 7, 96, fig. 3), PA 8796 (♀ RV: Pl. 7, 96, fig. 2), PA 8797 (♀ RV: Pl. 7, 94, fig. 1; Pl. 7, 96, fig. 1), PA 8798 (juv. - 1 RV: Pl. 7, 98, fig. 3; Pl. 7, 100, fig. 1), PA 8799 (juv. - 3 RV: Pl. 7, 100, fig. 3).

All figured material is from the type locality, between Elche de la Sierra and Ferez, Spain; lat. 38° 22' N, long. 02° 03' W; Miocene — Pliocene, yellowish calcarenite with *Chara* and abundant small gastropods; coll. by Dr. T. Geel.


*Diagnosis*: Medium-sized species of *Cyprideis* with box-like shape and conspicuous surface pitting. High nodding potential. Posteroventral node tends to be elongated and dorsoventrally compressed. Anomalous node may develop near posterodorsal corner. Anterior margins non-denticulate. Right valve with four posteroventral spines, some or all of which may be somewhat under-developed in the adults.

---

Explanation of Plate 7, 92

Fig. 1, ♂ LV, ext. lat. (paratype, PA 8784, 910 µm long); fig. 2, ♂ RV, ext. lat. (paratype, PA 8783, 880 µm long); fig. 3, ♂ car., ext. dors. (paratype, PA 8786, 890 µm long).  
Scale A (250 µm; x 68), figs. 1 - 3.











**Remarks:** *C. exuberans* is a very distinctive species that is not likely to be confused with any other described taxon of the genus. The nodding characteristics are quite exceptional in several respects. As many as seven nodes may develop on each valve but their location deviates from Sandberg's well-known scheme (see text-fig. 1): there is an anomalous node position in the posterodorsal area whereas Sandberg's position no. 5 is unoccupied (see P. A. Sandberg, *Stockh. Contr. Geol.*, 12, 42, 1964). Another notable feature is that the whole of the population seems to be capable of developing nodes: among the adults of the type sample (i.e. 249 left valves, 238 right valves and 30 carapaces) not a single smooth shell was found. A third anomaly is that the species does not comply with the rule that nodes are more common on juveniles than they are on adults (Sandberg, *op. cit.*, 47; see also T. I. Kilenyi, *Micropaleontology*, 18, 47 - 63, 1972). In *Cyprideis exuberans* sp. nov. the nodes make their ontological appearance in the juvenile - 2, in the type sample c. 30% of the individuals being affected. In the penultimate instar all but very few of the right valves are noded whereas c. 25% of the left valves are smooth.

To all appearance there is bimodality in the size distributions of the juv. - 1 and juv. - 2 instars (see text-figs. 2, 3). This is indicative of sexual dimorphism. As the same thing has been reported for *C. baetica* van Harten (D. van Harten, *Scripta Geologica*, 32, 6, 1975) and *C. torosa* (Jones) (W. - M. Rohr, *N. Jb. Geol. Paläont. Abh.* 158, 346 - 380, 1979), the existence of pre-adult sexual dimorphism may well be a generic feature in *Cyprideis*.

From the associated fauna and flora a shallow oligohaline, possibly even limnetic, palaeo-environment is inferred for the type sample.


#### Explanation of Plate 7, 94

Fig. 1, ♀ RV, int. lat. (PA 8797, 730  $\mu$ m long); fig. 2, ♀ LV, int. lat. (PA 8794, 780  $\mu$ m long); fig. 3, ♀ car., ext. vent. (paratype, PA 8785, 800  $\mu$ m long).

Scale A (250  $\mu$ m; x 75), figs. 1 - 3.

**Distribution:** So far, the species is only known from the type locality. There is some doubt about the actual age of the *stratum typicum*. According to L. Jerez-Mir (*Geología de la zona prebetica, en la transversal de Elche de la Sierra y sectores adyacentes*, thesis University of Granada, 750 pp., 1973) the lacustrine deposits at issue most probably belong to the Upper Miocene; however, a Pliocene age would not be utterly out of the question (Jerez-Mir, *op. cit.*, 505 - 508).

**Acknowledgement:** I am indebted to Tiny Geel for providing the material on which this paper is based and for supplying information on the stratigraphical setting.





Text-fig. 1. Ext. lat. view of ♀ RV showing possible positions of nodes in *Cyprideis exuberans* sp. nov. (A) as compared with Sandberg's general scheme (B). B after Sandberg (1964).

#### Explanation of Plate 7, 96

Fig. 1, ♀ RV, int. lat., musc. sc. (PA 8797); fig. 2, ♀ RV, ext. dors. (PA 8796, 780  $\mu$ m long); fig. 3, ♀ LV, ext. dors., ant. and median hinge elements (PA 8795); fig. 4, ♂ LV, ext. lat., detail of post. vent. area (PA 8789).

Scale A (50  $\mu$ m; x 375), fig. 1; scale B (250  $\mu$ m; x 75), fig. 2; scale C (100  $\mu$ m; x 190), figs. 3, 4.















Text-fig. 2. Length/height diagram for left valves in type sample.

## Explanation of Plate 7, 98

Fig. 1, juv. - 1 LV, ext. lat. (PA 8792, 700  $\mu$ m long); fig. 2, juv. - 1 LV, ext. lat. (PA 8791, 680  $\mu$ m long); fig. 3, juv. - 1 RV, int. obl., post. vent. spines (PA 8798).

Scale A (250  $\mu$ m; x 81), figs. 1, 2; scale B (50  $\mu$ m; x 407), fig. 3.




Text-fig. 3. Histogram of length for left valves in type sample.

## Explanation of Plate 7, 100

Fig. 1, juv. - 1 RV, int. lat. (PA 8798, 650  $\mu$ m long); fig. 2, juv. - 1 RV, ext. lat. (PA 8793, 640  $\mu$ m long); fig. 3, juv. - 3 RV, ext. lat. (PA 8799, 410  $\mu$ m long).

Scale A (250  $\mu$ m; x 81), figs. 1, 2; scale B (150  $\mu$ m; x 145), fig. 3.









ON *XESTOLEBERIS POSTANGULATA* BATE & SHEPPARD sp. nov.

by Raymond H. Bate & Lesley M. Sheppard  
(British Museum [Natural History], London)

*Xestoleberis postangulata* sp. nov.

*Holotype*: Brit. Mus. (Nat. Hist.) 1980.147, ♀ car.

[Paratypes: Brit. Mus. (Nat. Hist.) 1980.148 - 154]

*Type locality*: Abu Dhabi lagoon, central lagoon terrace, sample no. 6514b.

*Derivation of name*: Pertaining to the distinct posteroventral projection on each valve.

*Figured specimens*: Brit. Mus. (Nat. Hist.) nos. 1980.147 (holotype, ♀ car: Pl. 7, 102, fig. 1), 1980.148 (♀ RV: Pl. 7, 102, fig. 2), 1980.149 (♀ LV: Pl. 7, 102, fig. 3; Pl. 7, 104, figs. 5, 6), 1980.150 (♀ car: Pl. 7, 104, fig. 1), 1980.151 (♀ car: Pl. 7, 104, figs. 2, 9), 1980.152 (♀ RV: Pl. 7, 104, figs. 3, 4, 8), 1980.153 (LV: Pl. 7, 104, fig. 7).

*Diagnosis*: Small oval *Xestoleberis* having sharply angled posteroventral corner. Hinge paraperatodont; sieve plates with central setal pore and two or three concentric rows of sieve pores. Anterior and posterior marginal canals branching.

Explanation of Plate 7, 102

Fig. 1, ♀ car. lt. lat. (holotype, 1980. 147, 417 µm long); fig. 2, ♀ RV, ext. lat. (paratype, 1980.148, 417 µm long); fig. 3, ♀ LV, int. lat. (paratype, 1980.149, 383 µm long).

Scale A (100 µm; x 143), figs. 1, 2; scale B (100 µm; x 151), fig. 3.

*Remarks*: *X. postangulata* is a rare member of the littoral phytal fauna of the Abu Dhabi lagoon and nearshore shelf. In the lagoon, the species is present within the deeper water channels (depth about 7m), near the entrance to the lagoon and on the shallower water terraces (see text-fig. 3).

The species is dimorphic, dimorphism being recognised by the more slender outline of the males in dorsal view (♂ paratype 1980.10 measuring length 417 µm; height 255 µm; width 221 µm.) By comparison the ♀ holotype 1980.1 measures: length 417 µm; height 262 µm; width 245 µm. The hinge in *X. postangulata* is paraperatodont, a term introduced by Bate, 1972 (*Spec. pap. Palaeont.*, 10, 45) for certain members of the Cytheruridae; this type of hinge being characterised by having strongly developed dentition at the anterior and posterior ends of a smooth median element (see Pl. 7, 104, figs. 5, 6).



In addition to the small sieve plates present on the lateral and ventral surfaces, a smaller number of simple, rimmed, normal pores are present (not figured here). Apart from the present species only *X. ventribullata* Hartmann, 1962 (*Mitt. Hamburg Zool. Mus. Inst.*, 60, 227) from the Atlantic Coast of Argentina has a comparable posteroventral angular projection of the carapace but this is not so well developed as in *X. postangulata*. There is also a marked difference in the anterior margin of both species, that of *X. ventribullata* being very much broader than in *X. postangulata* (see text-fig. 2), while the anterior marginal canals are totally dissimilar.

Explanation of Plate 7, 104

Fig. 1, ♀ car., ext. dors. (paratype, 1980.150, 408 µm long); fig. 2, ♀ car., ext. vent. (paratype, 1980.151, 425 µm long); figs. 3, 6, ♀ RV, hinge (paratype, 1980.152); figs. 4, 5, ♀ LV, hinge (paratype, 1980.149); fig. 7, ♀ LV, RPC (paratype, 1980.153); fig. 8, ♀ RV, int. lat. (paratype, 1980.152, 417 µm long); fig. 9, ♀ car. NPC sieve pl. (paratype, 1980.151).

Scale A (100 µm; x 98), fig. 1; scale B (100 µm; x 94), fig. 2; scale C (200 µm; x 321), figs. 3, 4, 5, 6; scale D (200 µm; x 236), fig. 7; scale E (100 µm; x 95), fig. 8; scale F (25 µm; x 1190), fig. 9.









Text-fig. 1. Muscle scars, LV, paratype 1980.149 x 700.



Text-fig. 2. Comparison of LV anterior outlines of:  
A - *X. ventribullata* Hartmann 1962.  
B - *X. postangulata* sp. nov.



Text-fig. 3. Distribution of *Xestoleberis postangulata* sp. nov. within and outside Abu Dhabi lagoon. Arrows point to location of sample of which only those indicated, out of a total of 83, contained the species.







ON *QUADRALEBERIS EXQUISITA* BATE & SHEPPARD gen. et sp. nov.

by Raymond H. Bate & Lesley M. Sheppard  
(British Museum [Natural History], London)

Genus *QUADRALEBERIS* gen. nov.

Type species: *Quadraleberis exquisita* sp. nov.

Gender: Feminine.

**Derivation of name:** Latin *quadrus*, square – relating to the very square outline of the carapace + *leberis*, used here to relate the genus to the family Trachyleberideidae and not to the Latin word for skin.

**Diagnosis:** Genus of Trachyleberideidae having quadrate (female) to rectangular (male) outline, dorsal and ventral margins straight and almost parallel. Eye node distinct. External muscle scar node absent. Carapace triangular in dorsal view, widening to posterior. Anterior and posterior margins spinose. Posterior ventro-lateral region with short but distinct projection. Hinge holamphidont. V-shaped frontal scar, undivided adductors. Normal pore apertures simple; anterior marginal pore canals long, slender, numerous; duplicature broad, without a vestibule. LV larger than RV.

Explanation of Plate 7, 108

Fig. 1, ♂ LV, ext. lat. (holotype, 1980.155, 950  $\mu$ m long); fig. 2, ♂ RV, ext. lat. (paratype, 1980.156, 900  $\mu$ m long); fig. 3, ♀ RV, ext. lat. (paratype, 1980.157, 880  $\mu$ m long).

Scale A (200  $\mu$ m; x 63), fig. 1; scale B (200  $\mu$ m; x 67), figs. 2, 3.

**Remarks:** *Quadraleberis*, although lacking an external muscle scar node typical of the Trachyleberideidae is, nevertheless, considered to belong to this family on all the other characters of the carapace. *Quadracythere* Hornibrook, 1952 has a divided frontal scar as well as a ventrolateral keel and is easily distinguished on this evidence (as also is the subgenus *Hornibrookella* Moos, 1965). Similarly, *Bensoncythere* Hazel, 1967 has a divided frontal scar and thus belongs to the Hemicytheridae. *Robertsonites* Swain, 1963 is a trachyleberid genus having a quadrate outline but that is perhaps the only similarity when compared with *Quadraleberis*. Although there are a number of genera that superficially appear to have features in common with *Quadraleberis* this is not borne out on closer examination. Indeed, it is considered that our new genus most probably represents a new subfamily of the Trachyleberideidae characterised by the absence of an external muscle scar node.

*Quadraleberis exquisita* sp. nov.

1976 *Quadracythere* (*Quadracythere*) sp. A Kwang Ho Paik, *Doctorate Diss. Univ. Kiel*, 85, pl. 6, figs. 101 - 104.

1977 *Quadracythere* (*Quadracythere*) sp. A Kwang Ho Paik, "Meteor" *Forsch-Ergebnisse*, C, 28, pl. 6, figs. 101 - 104.

**Holotype:** Brit. Mus. (Nat. Hist.) no. 1980.155, ♂ LV.  
[Paratypes; Brit. Mus. (Nat. Hist.) nos. 1980.156 - 161].

**Type locality:** Sample G3 station 2, Persian Gulf (see text-fig. 2).



**Figured specimens:** Brit. Mus. (Nat. Hist.) nos. 1980.155 (holotype, ♂ LV: Pl. 7, 108, fig. 1; Pl. 7, 110, fig. 2), 1980.156 (♂ RV: Pl. 7, 108, fig. 2), 1980.157 (♀ RV: Pl. 7, 110, fig. 7), 1980.158 (♂ RV: Pl. 7, 110, figs. 1, 3, 4), 1980.159 (♀ RV: Pl. 7, 108, fig. 3), 1980.160 (juv. RV: Pl. 7, 110, fig. 6), 1980.161 (♀ RV: Text-fig. 1).

Explanation of Plate 7, 110

Fig. 1, musc. sc. (paratype, 1980.158); fig. 2, ornament (holotype, 1980.155); figs. 3, 4, ♂ RV, terminal hinge elements (paratype, 1980.158); fig. 5, ♀ LV, ext. lat. (specimen lost); fig. 6, juv. RV, ext. lat. (paratype, 1980.160, 680  $\mu$ m long); fig. 7, ♀ RV, dors. (paratype, 1980.157, 825  $\mu$ m long).

Scale A (20  $\mu$ m; x 550), fig. 1; scale B (50  $\mu$ m; x 211), fig. 2; scale C (100  $\mu$ m; x 51), figs. 3, 4; scale D (200  $\mu$ m; x 51), fig. 5; scale E (200  $\mu$ m; x 66), fig. 6; scale F (100  $\mu$ m; x 109), fig. 7.











**Diagnosis:** *Quadraleberis* with a striking trefoil ornamentation arranged anteriorly in rows paralleling anterior margin, elsewhere more irregular with some tendency towards being concentrically aligned. Intensity of ornamentation greater in juvenile instars. Hinge holamphidont with dentate/loculate median element. Up to 46 long, slender, anterior marginal canals.

**Remarks:** *Quadraleberis exquisita* sp. nov. has the same trefoil ornamentation as described for *Anebocythereis amoena* Bate, 1972 (*Spec. pap. Paleont.*, 10, 53, pl. 14, figs. 1, 2) but differs generically on carapace outline. Indeed the very angular outline of this species clearly sets it apart from all other members of the Trachyleberideidae. The muscle scars (Pl. 7, 110, fig. 1) are typical of this family though the absence of a muscle scar pit (and external node) are in contrast with members of the Trachyleberideinae. This species has been described by Paik 1976 from the deeper waters of the Persian Gulf with sizes ranging between 0.80 - 0.82mm for the female and 0.86 - 0.90mm for the male. The length of our specimens is slightly larger than this with, at the same time, a slightly increased number of anterior marginal canals (up to 46 instead of 39 - 42). There is no doubt, however, that our species is the same as that described by Paik.

**Distribution:** Recorded here from stations G6 st6 (lat. 26° 35' N, long. 53° 24' E) and G3 st2 (lat. 26° 10' N, long. 55° 20' E) in the deeper waters of the Persian Gulf from material collected by Dr Graham Evans, Imperial College, London in 1962. Paik's material, collected during the "Meteor"-Expedition in 1965, also comes from the deeper water of the Gulf, as well as from the Gulf of Oman, from depths ranging from 63 to 196 metres and from a calc. marl substrate. Shallow water faunas described by Jain, 1978 (*Bull. Ind. Geol. Assoc.*, 11) from the West Coast of India do not contain this species even though many other species are common to the two regions. Water depth is thus an important factor in the distribution of *Q. exquisita*.



Text-fig. 1, anterior marginal pore canals, ♀ RV. 1980.161 x 148.



Text-fig. 2, distribution of *Quadraleberis exquisita* gen. et sp. nov. in the deeper water regions of the Persian Gulf. Type locality, sample G3 st. 2, ringed.







**ON *MICROPNEUMATOCY THERE TUMIDA* STEPHENS & WARE sp. nov.**

by Jill Stephens and Martin Ware  
(*Soekor, Johannesburg; University College of Wales, Aberystwyth*)

*Micropneumatocythere tumida* sp. nov.

**Holotype:** Brit. Mus. (Nat. Hist.) **OS 11739**, ♀ LV.  
[Paratypes: Brit. Mus. (Nat. Hist.) nos. **OS 11740 - 11746**].

**Type locality:** Kirtlington Mammal Bed (see E. F. Freeman, *Palaeontology*, **22** (1), 135 - 166, 1979), Forest Marble, Upper Bathonian, Old Cement Works quarry, Kirtlington, Oxfordshire. Nat. Grid Ref.: SP 494199.

**Derivation of name:** Latin *tumidus*, swollen: refers to ventrolateral tumidity.

**Figured specimens:** Brit. Mus. (Nat. Hist.) nos. **OS 11739** (holotype, ♀ LV: Pl. 7, 114, fig. 1; Pl. 7, 116, fig. 4), **OS 11740** (♀ RV: Pl. 7, 114, fig. 2), **OS 11741** (♂ LV: Pl. 7, 114, fig. 3), **OS 11742** (♂ RV: Pl. 7, 114, fig. 4; Pl. 7, 116, fig. 5), **OS 11743** (♂ car.: Pl. 7, 116, fig. 1), **OS 11744** (♀ LV: Pl. 7, 116, fig. 2), **OS 11745** (♂ RV: Pl. 7, 116, fig. 3), **OS 11746** (♂ RV: Pl. 7, 116, fig. 6). All from the type locality and horizon.

**Diagnosis:** An inflated, dimorphic species of *Micropneumatocythere*, suboval in lateral outline, with maximum tumidity at midlength ventrolaterally and strong ventral overlap. Cardinal angles indistinct. Surface ornamented with faint open reticulation and fine punctae. Normal pore canals contain sieve-plates. Other features as for genus.

**Explanation of Plate 7, 114**

Fig. 1, ♀ LV, ext. lat. (holotype, **OS 11739**, 530 µm long); fig. 2, ♀ RV, ext. lat. (paratype, **OS 11740**, 470 µm long); fig. 3, ♂ LV, ext. lat. (paratype, **OS 11741**, 580 µm long); fig. 4, ♂ RV, ext. lat. (paratype, **OS 11742**, 570 µm long).  
Scale A (200 µm; x 82), fig. 1; scale B (200 µm; x 86), fig. 2; scale C (200 µm; x 81), fig. 3; scale D (200 µm; x 88), fig. 4.

Size:	Range (Mean):	♀♀ LV	♀♀ RV	♂♂ LV	♂♂ RV
	Length µm (10 valves)	470 - 550 (501)	460 - 510 (479)	560 - 580 (575)	550 - 600 (567)
	Height µm (10 valves)	340 - 390 (359)	300 - 330 (313)	340 - 370 (356)	300 - 350 (323)

**Remarks:** At the type horizon *Micropneumatocythere tumida* was associated with *M. brendae* Sheppard, *M. postrotunda* Bate, *M. subconcentrica* (Jones), and *M. cracens* Bate & Sheppard. As with *M. cracens*, males of *M. tumida* are the more distinctive, readily distinguished from all known members of the genus by their midlength tumidity. Females are very similar to those of *M. brendae* and *M. falcata* Sheppard. They are distinguished from smooth forms of *M. brendae* by their characteristic tumidity, a less strongly arched LV dorsal outline, and absence of vertical ridges in the accommodation groove. They have a shorter caudal process than *M. falcata*, the LV posterodorsal slope is convex rather than slightly concave, and maximum tumidity is more ventral.

Sieve-plates have not previously been recorded in this genus. *M. tumida* also exhibits precocious sexual dimorphism.



Because of the mixed assemblages from which *M. tumida* has been recovered, its salinity tolerance is uncertain: it was probably a shallow-marine species with only limited tolerance of brackish conditions.

**Distribution:** *M. tumida* is known only from the type locality and the White Limestone and Forest Marble of the neighbouring Portland Cement Manufacturers' quarry at Shipton-on-Cherwell (Nat. Grid Ref.: SP 477175).

**Explanation of Plate 7, 116**

Fig. 1, ♂ car., ext. dors. (paratype, **OS 11743**, 560 µm long); fig. 2, ♀ LV, int. lat. (paratype, **OS 11744**, 550 µm long); fig. 3, ♂ RV, hinge (paratype, **OS 11745**, 570 µm long); fig. 4, ♀ LV, ornament and sieve-pores (holotype, **OS 11739**, 530 µm long); fig. 5, ♂ RV, sieve-plate (paratype, **OS 11742**, 570 µm long); fig. 6, ♂ RV, adductor scars (paratype, **OS 11746**, 570 µm long).  
Scale A (200 µm; x 83), fig. 1; scale B (200 µm; x 83), fig. 2; scale C (100 µm; x 134), fig. 3; scale D (40 µm; x 360), fig. 4; scale E (4 µm; x 1,813), fig. 5; scale F (40 µm; x 285), fig. 6.











ON *PROCYTHEROPTERON PROLONGATUM* (SHARAPOVA)

by Nicholas Fuller  
(University College, London)

*Procytheropteron prolongatum* (Sharapova, 1939)

- 1939 *Cytheropteron prolongatum* sp. nov. E. G. Sharapova, *Trudy NGRI*, A126, 23, pl. 2, fig. 23.  
1955 *Procythere prolongata* (Sharapova, 1939); P. S. Lyubimova, *Trudy VNIGRI*, new series, 84, 112, pl. 8, figs. 2a - c.  
non v. 1955 *Cytheropteron* cf. *prolongatum* (Sharapova, 1939); G. Schmidt, *Abh. senckenb. naturf. Ges.*, 491, 60, pl. 2, fig. 34.  
1978 *Procytheridea prolongata* (Sharapova, 1939); D. M. Pyatkova & M. N. Permyakova, *Jurassic Foraminifera and Ostracoda of the Ukraine*, 154, pl. 69, figs. 5a - d.

**Holotype:** Collection of the Petroleum Geological Prospecting Institute (NGRI) Leningrad, no. 51 - 20.

**Type locality:** Depth 267 - 273m in Well no. 1, 1km E of the Lenin Farm, left bank of Chalykla River, along flank of the Ozinki Dome, in Western Obshchiy Syrt, Soviet Russia. Lower Volga stage, *T. Perisphinctes panderi* zone.

---

Explanation of Plate 7, 118

Fig. 1, ♀ LV, ext. lat. (OS 11767, 520  $\mu$ m long); fig. 2, ♀ LV, int. lat. (OS 11768, 520  $\mu$ m long); fig. 3, ♀ RV, ext. lat. (OS 11769, 510  $\mu$ m long).

Scale A (100  $\mu$ m; x 90), fig. 1; scale B (100  $\mu$ m; x 100), fig. 2; scale C (100  $\mu$ m; x 105), fig. 3.

**Figured specimens:** Brit. Mus. (Nat. Hist.) nos. OS 11767 (♀ LV: Pl. 7, 118, fig. 1), OS 11768 (♀ LV: Pl. 7, 118, fig. 2; Pl. 7, 124, fig. 1), OS 11769 (♀ RV: Pl. 7, 118, fig. 3), OS 11770 (♂ LV: Pl. 7, 120, fig. 1), OS 11771 (♀ RV: Pl. 7, 120, fig. 2; Pl. 7, 124, fig. 3), OS 11772 (♂ RV: Pl. 7, 120, fig. 3), OS 11773 (juv. A - 1 LV: Pl. 7, 122, fig. 1; Pl. 7, 124, fig. 5), OS 11774 (juv. A - 2 LV: Pl. 7, 122, fig. 2), OS 11775 (juv. A - 3 LV: Pl. 7, 122, fig. 3), OS 11776 (♀ LV: Pl. 7, 124, fig. 2), OS 11777 (♀ RV: Pl. 7, 124, fig. 4).



Specimens OS 11773, OS 11776 from Bed 8 (*Subplanites pseudoscythicus* Zone, Lower Volgian), others from Beds 3 and 4 (*Aulacostephanus autissiodorensis* Zone, Upper Kimmeridgian) at Gorodische, 25km north of Ul'yanovsk, River Volga, U.S.S.R.. Bed numbers of Mesezhnikov, M. S., Dain, L. G., Kuznetsova, K. I. and Yakovleva, S. P., *International Colloquium on Upper Jurassic stratigraphy and the Jurassic/Cretaceous boundary in the Boreal Realm - Jurassic/Cretaceous boundary beds in the Middle Volga area (A prospectus to Geological Excursions)*, All-Union Petroleum Research Geological Prospecting Institute, (VNIGRI), Leningrad, fig. 1, 1977.

---

Explanation of Plate 7, 120

Fig. 1, ♂ LV, ext. lat. (OS 11770, 545  $\mu$ m long); fig. 2, ♀ RV, int. lat. (OS 11771, 490  $\mu$ m long); fig. 3, ♂ RV, ext. lat. (OS 11772, 565  $\mu$ m long).

Scale A (100  $\mu$ m; x 100), fig. 1; scale B (100  $\mu$ m; x 110), fig. 2; scale C (100  $\mu$ m; x 95), fig. 3.











**Diagnosis:** Rectangular, inflated, tapering to prominent caudal process. Coarsely reticulate surface ornament. Narrow anterior marginal rim. Some sieve type normal pores, particularly in posterodorsal portion of valves. Anterior pore canals vary in number between 7 and 9, simple and relatively straight, widely spaced, not all reach margin. No pore canals observed in posterior region. Pronounced sexual dimorphism. Hinge hemimerodont, in the LV terminal crenulate sockets are separated by a smooth median bar. Adults and three juvenile stages recognised.

**Remarks:** First assigned to the genus *Cytheropteron* Sars 1866 in 1939 by E. G. Sharapova, this ostracod lacks true *Cytheropteron* characters in having a smooth median hinge bar and no prominent posteroventral alar processes. Subsequently assigned to the genus *Protocythere* Triebel 1938 by P. S. Lyubimova (1955), this species is not considered to belong to the genus through lack of an antimerodont hinge and 3 characteristic longitudinal ridges of *Protocythere*. More recently D. M. Pyatkova and M. N. Permyakova (1978) attributed *P. prolongatum* to *Procytheridea* Peterson 1954, however this genus is also characterised by an antimerodont hinge, possesses wide marginal areas and has a different central muscle scar pattern. Some confusion also arises in assigning this species to the genus *Procytheropteron* as this genus has been described twice, first by P. S. Lyubimova (1955), whose description complies with *P. prolongatum* in all respects saving that she describes the radial pore canals as "frequent and present in great numbers at the anterior end". In 1956 M. I. Mandelstam also described a genus *Procytheropteron* and *P. prolongatum* agrees more closely with his description, differing in one small detail, namely that Mandelstam's genus has a gently convex LV dorsal margin. This difference is not considered significant here at a generic level.

---

Explanation of Plate 7, 122

Fig. 1, juv. A - 1, LV, ext. lat. (OS 11773, 430  $\mu$ m long); fig. 2, juv. A - 2, LV, ext. lat. (OS 11774, 415  $\mu$ m long); fig. 3, juv. A - 3, LV, ext. lat. (OS 11775, 340  $\mu$ m long).

Scale A (100  $\mu$ m; x 116), fig. 1; scale B (100  $\mu$ m; x 120), fig. 2; scale C (100  $\mu$ m; x 147), fig. 3.

**Remarks:** Van Morkhoven (*Post-Palaeozoic Ostracoda*, 2, 258 and 385, Elsevier, 1963) considers *Procytheropteron* (cont'd.) Lyubimova a synonym of *Neocythere* Mertens 1956. *Neocythere* s.s., however, possesses an amphidont hinge. The subgenus *N. (Physocythere)* Kaye 1963 has a merodont hinge, but lacks other features of *Procytheropteron*, notably the well developed caudal process. A caudal process is not specified in either diagnosis but is a feature of *P. obesum* Lyubimova 1955, the type-species of *Procytheropteron*. One feature of *P. prolongatum* not hitherto mentioned in any descriptions of this species is the presence of several well developed, though small "sieve-type" normal pores in the posterodorsal region (see Pl. 7, 124, fig. 5).

*P. prolongatum* is found in association with *Galliaecytheridea miranda* (Lyubimova), *G. volgaensis* (Mandelstam), *G. mandelstami* (Lyubimova), *G. gorodischensis* (Fuller and Lord), *G. ramosa* (Lyubimova), *Protocythere bisulcata* (Sharapova), *Mandelstamia ventrocornuta* (Sharapova), *Oligocythereis kostytschevkaensis* (Lyubimova) and *Cytherella* spp.

**Distribution:** Upper Kimmeridgian, Lower and Middle Volgian of Gorodische in the Volga region; known also from the Ukraine. A similar form is recorded by G. Schmidt (1955, *op. cit.*) from 30km S of Hannover.



---

Explanation of Plate 7, 124

Fig. 1, ♀ LV, int. lat. musc. sc. (OS 11768); fig. 2, ♀ LV, int. lat. hinge (OS 11776); fig. 3, ♀ RV, int. lat. hinge (OS 11771); fig. 4, ♀ LV, ext. lat. detail of ornament (OS 11777); fig. 5, juv. A - 1, ext. lat. sieve pore (OS 11773).

Scale A (100  $\mu$ m; x 345), fig. 1; scale B (100  $\mu$ m; x 195), fig. 2; scale C (40  $\mu$ m; x 190), fig. 3; scale D (20  $\mu$ m; x 370), fig. 4; scale E (4  $\mu$ m; x 3700), fig. 5.











**ON SEMICYTHERURA TELA HORNE & WHITTAKER sp. nov.**

by David J. Horne and John E. Whittaker

(Geology Department, Bristol University, and British Museum [Natural History], London)

*Semicytherura tela* sp. nov.

1969 *Semicytherura nigrescens* (Baird) pars; I. Yassini, *Bull. Inst. Geol. Bassin Aquitaine*, 7, 88, pl. 17.

**Holotype:** Brit. Mus. (Nat. Hist.) 1980.134, ♀ LV.

[Paratypes: Brit. Mus. (Nat. Hist.) nos. 1980.135 - 139].

**Type locality:** Blue Anchor, Somerset, W England (S shore of Severn Estuary); approx. lat. 51° 12' N, long. 3° 24' W. Recent, littoral.

**Derivation of name:** Latin, *tela*, *ae* — woven stuff or web. The fine ornamentation gives the carapace the appearance of being draped with cobwebs or fine netting.

**Explanation of Plate 7, 126**

Fig. 1, ♀ LV, ext. lat. (holotype, 1980.134, 360 µm long); fig. 2, ♂ LV, ext. lat. (paratype, 1980.135, 380 µm long); fig. 3, juv. -1 LV, ext. lat. (paratype, 1980.136, 330 µm long).

Scale A (100 µm; x 170), figs. 1 - 3.

**Figured specimens:** Brit. Mus. (Nat. Hist.) nos. 1980.134 (holotype, ♀ LV: Pl. 7, 126, fig. 1), 1980.135 (♂ LV: Pl. 7, 126, fig. 2; Pl. 7, 128, fig. 3), 1980.136 (juv. -1 LV: Pl. 7, 126, fig. 3), 1980.137 (♀ LV: Pl. 7, 128, fig. 1), 1980.138 (♂ car.: Pl. 7, 128, fig. 2), 1980.139 (♂ copulatory appendage: Text-fig. 1).

1980.134 - 137, 139, collected alive from *Corallina officinalis* in rocks-pools at the type locality by D. J. Horne, 22nd July 1978; salinity and water temperature ranged between 28.5 and 29.2‰, and 10 to 19°C, respectively, measured over a spring tide. 1980.138 was collected alive from *Cladophora* by J. E. Whittaker, at Osmington Mills, Weymouth Bay, S England (approx. lat. 50° 38' N, long 2° 23' W), 3rd August 1968; salinity 33.9‰, water temperature 19°C.


**Diagnosis:** Very small *Semicytherura* (<0.40mm long) with ornamentation of fine lateral ridges and associated rows of shallow pits, developing into reticulation in the posterior half. Males smaller, more elongate and inflated posteriorly. Copulatory organ characteristic, with a triangular lappet and a finger-like projection on the ventral margin.

**Explanation of Plate 7, 128**


Fig. 1, ♀ LV, int. lat. (paratype, 1980.137, 380 µm long); fig. 2, ♂ car., ext. dors. (paratype, 1980.138, 370 µm long); fig. 3, ♂ LV, detail of mid-posterior region (paratype, 1980.135).

Scale A (100 µm; x 170), figs. 1, 2; scale B (10 µm; x 1,100). fig. 3.









Text-fig. 1, ♂ rt. copulatory appendages (1980.139); text-fig. 2, ♂ rt. copulatory appendage (1980.140) of *Semicytherura nigrescens* (Baird), shown for comparative purposes. Both come from specimens collected at Blue Anchor, Severn Estuary, from littoral algae.

*Remarks:* This species has probably been confused in the past with *Semicytherura nigrescens* (Baird) (see Whittaker, *StereAtlas of Ostracod Shells*, 2, 69 - 76 1974), from which it can be distinguished by its smaller size, stronger ornamentation, and distinctive copulatory organ (cf. text-figs. 1 and 2). *S. tela* was found by one of us (J. E. W.) in the "type-material" of *S. nigrescens sensu* Yassini, from the Arcachon Basin, at Bordeaux University, and would seem to possess a more extensive geographical distribution than is at present recognised. It is much too small to be considered synonymous with *S. similis* (Sars, 1866).

*Distribution:* Recent; *S. tela* has so far been recorded from three regions on the British coasts. In the Fleet and Weymouth Bay, S England, it was found on filamentous green-algae such as *Enteromorpha* and *Cladophora*, in salinities from 31 - 35‰, while in the Bristol Channel and Severn Estuary, it has been collected at several localities between Porlock Bay and Sand Point, in intertidal samples of *Corallina*, *Laminaria* holdfasts and *Sabellaria* worm-tubes, within the salinity range 20 - 35‰. K. Trier has recently sent us material collected from tidal pools in S W Wales. As mentioned above it also occurs in the Arcachon Basin, S W France.







ON *ACANTHOCYHEREIS DECORIS* SIDDIQUI

by Qadeer A. Siddiqui  
(Saint Mary's University, Halifax, Canada)

*Acanthocythereis decoris* Siddiqui, 1971

1971 *Trachyleberis (Acanthocythereis) decoris* sp. nov. Q. A. Siddiqui, *Bull. Brit. Mus. nat. Hist. (Geol.) Suppl.* 9, 83, pl. 42, figs. 3 - 6, 8, 9.

*Holotype*: Brit. Mus. (Nat. Hist.) IO 4359, ♂ car.

*Type locality*: Rakhi Nala section, Sulaiman Range, Pakistan; lat. 29° 57.5' N, long. 70° 60' E. Upper Chocolate Clays, Kirthar Formation, Upper Eocene.

*Figured specimens*: Brit. Mus. (Nat. Hist.) nos. OS 10040 (♀ car.: Pl. 7, 132, fig. 1), OS 11737 (♂ car.: Pl. 7, 132, figs. 2, 3; Pl. 7, 134, figs. 1 - 3), both from the type locality above; Upper Chocolate Clays, sample no. 3604, Kirthar Formation, Middle Eocene.

---

Explanation of Plate 7, 132

Fig. 1, ♀ car., ext. lt. lat. (OS 10040, 765 µm long); fig. 2, ♂ car., ext. rt. lat. (OS 11737, 791 µm long); fig. 3, ♂ car., ext. post. (OS 11737).

Scale A (200 µm; x 77), fig. 1; scale B (200 µm; x 82), fig. 2; scale C (100 µm; x 96), fig. 3.

*Diagnosis*: A species of *Acanthocythereis* in which the eye-tubercle is rounded and distinct; subcentral tubercle indistinct; shell surface reticulate with conjunctive papillae; anterior, posterior and ventral margin decorated with a double row of short spines and/or papillae.

*Remarks*: The subcentral tubercle in this species seems to vary; in some specimens it is almost indistinct. In a few specimens papillose spines occur on shell surface.

*Distribution*: Rakhi Nala and Zao River sections of the Sulaiman Range, Pakistan (Middle — Upper Eocene).



---

Explanation of Plate 7, 134

Fig. 1, ♂ car., ext. dors. (OS 11737, 791 µm long); fig. 2, ♂ car., ext. vent. (OS 11737); fig. 3, ♂ car., ext. ant. (OS 11737).

Scale A (200 µm; x 79), fig. 1; scale B (200 µm; x 82), fig. 2; scale C (100 µm; x 96), fig. 3.











ON *CALLISTOCY THERE MEDITERRANEA* (MÜLLER)

by Neriman Doruk  
(Ege University, Izmir, Turkey)

*Callistocythere mediterranea* (Müller, 1894)

1894 *Cythere mediterranea* sp. nov. G. Müller, *Zool. stat. Naples, Monogr.*, no. 21, 353, pl. 27, fig. 34; pl. 28, fig. 13; pl. 32, fig. 34.

*Holotype*: Zoological Museum, Berlin (see Diebel, *Geologie*, 11, no. 2, 249, 1962).

*Type locality*: Bay of Naples, Italy. Recent.

*Figured specimens*: Brit. Mus. (Nat. Hist.) nos. **Io 5175** (♂ RV: Pl. 7, 136, fig. 1), **Io 5176** (♀ LV: Pl. 7, 136, fig. 2; Pl. 7, 138, figs. 1, 3), **Io 5177** (♂ RV: Pl. 7, 138, fig. 2). **Io 5175** was dredged from Urla Bay, west coast of Turkey and is Subrecent, shallow water marine (littoral). Approx. lat. 38° 19' N, long. 26° 47' E. **Io 5176** and **Io 5177** were from drillings off the south coast of Turkey, 450ft. below sea floor, Plio-Pleistocene, presumed shallow water marine (littoral); approx. lat. 36° 26' N, long. 35° 04' E.

## Explanation of Plate 7, 136

Fig. 1, ♂ RV, ext. lat. (**Io 5175**, 512 µm long); fig. 2, ♀ LV, ext. lat. (**Io 5176**, 532 µm long).  
Scale A (250 µm; x 166), fig. 1; scale B (250 µm; x 156), fig. 2.



*Diagnosis*: Diagnostic rib pattern associated with nine anterior marginal denticulations.

*Remarks*: *Callistocythere mediterranea* has not been rediscovered from the type locality since G. W. Müller's first record. It seems, however, to be a common species in the Eastern Mediterranean.

*Distribution*: Recent; Gulf of Naples and Urla Bay (Turkey); Pleistocene, south coast of Turkey.

## Explanation of Plate 7, 138

Figs. 1, 3, ♀ LV, (**Io 5176**, 532 µm long): fig. 1, int. lat.; fig. 3, musc. sc.; fig. 2, ♂ RV, int. lat. (**Io 5177**, 530 µm long).  
Scale A (250 µm; x 126), fig. 1; scale B (250 µm; x 130), fig. 2; scale C (100 µm; x 453), fig. 3.











ON *CALLISTOCY THERE MONTANA* DORUK sp. nov.

by Neriman Doruk  
(Ege University, Izmir, Turkey)

*Callistocythere montana* sp. nov.

*Holotype*: Brit. Mus. (Nat. Hist.) no. Io 5168, ♂ RV.

*Type locality*: Road cutting 3km S W of Kuzucubelen (Mersin region of Turkey), approx. lat. 36° 48' N, long. 34° 27' E. Tortonian (Middle to Upper Miocene). Grey sandy clay with abundant foraminifera and molluscan shell fragments. Presumed intermediate depth — marine shelf.

*Derivation of name*: Latin “mountainous”, referring to ornamentation.

*Figured specimens*: Brit. Mus. (Nat. Hist.) nos. Io 5168 (♂ RV: Pl. 7, 140, fig. 1; Pl. 7, 142, figs. 2, 3), Io 5169 (♂ LV: Pl. 7, 140, fig. 2). All figured specimens from the type locality, about 4 metres from the base of the section.

---

Explanation of Plate 7, 140

Fig. 1, ♂ RV, ext. lat. (holotype, Io 5168, 462 µm long); fig. 2, ♂ LV, ext. lat. (Io 5169, 420 µm long).  
Scale A (250 µm; x 186), fig. 1; scale B (250 µm; x 205), fig. 2.

*Diagnosis*: Ornamentation diagnostic; surface between costae and tubercles smooth.

*Remarks*: Strength of costae and tubercles varies in intensity on the carapace. Sexual dimorphism pronounced; males less high posteriorly, females subrectangular. Ornamentation more strongly marked in males.



*Distribution*: Known, so far, only from the type locality.

---

Explanation of Plate 7, 142

Fig. 1, ♂ LV, int. lat. (specimen lost); figs. 2, 3, ♂ RV, (holotype Io 5168, 462 µm long); fig. 2, int. lat.; fig. 3, musc. sc.  
Scale A (250 µm; x 169), fig. 1; scale B (250 µm; x 151), fig. 2; scale C (100 µm; x 453), fig. 3.











ON *PARACYTHERIDEA INSCITA* DORUK sp. nov.

by Neriman Doruk  
(Ege University, Izmir, Turkey)

*Paracytheridea inscita* sp. nov.

*Holotype*: Brit. Mus. (Nat. Hist.) no. **Io 5209**, ♀ RV.

*Type locality*: Road cutting between Babatorun and Com, 1km S W of Babatorun (Turkey); approx. lat. 36° 04' N, long. 36° 15' E. Upper Miocene: yellow sandstone with abundant foraminifera and mollusc shells; presumed shallow water marine.

*Derivation of name*: Latin, "clumsy", referring to heavy ornamentation.

*Figured specimens*: Brit. Mus. (Nat. Hist.) nos. **Io 5209** (♀ RV: Pl. 7, 144, figs. 1, 3; Pl. 7, 146, figs. 2, 3), **Io 5210** (♂ LV: Pl. 7, 144, fig. 2), **Io 5211** (♂ LV: Pl. 7, 146, fig. 1). **Io 5209** and **Io 5211** are from the type locality, 10 metres from the base of the section. **Io 5210** is from a stream cutting 200 metres south of Sarıh, Turkey, 2 - 3 metres from the base of the section. Tortonian, bioclastic limestone with molluscan shells, presumed shallow water marine. Approx. lat. 36° 07' N, long. 36° 13' E.

Explanation of Plate 7, 144

Figs. 1, 3, ♀ RV (holotype, **Io 5209**, 678 µm long): fig. 1, ext. lat.; fig. 3, papillate sola; fig. 2, ♂ LV, ext. lat. (**Io 5210**, 673 µm long).

Scale A (250 µm; x 115), fig. 1; scale B (250 µm; x 113), fig. 2; scale C (50 µm; x 368), fig. 3.

*Diagnosis*: Heavily ornamented.

*Remarks*: Similar to *P. triquetra* (Reuss) but less triangular in lateral outline. Slightly variable in the strength of the ribs.



Males a little less high (see Pl. 7, 144, fig. 2) than females.

*Distribution*: Upper Miocene; Antakya region of Turkey.

Explanation of Plate 7, 146

Fig. 1, ♂ LV, int. lat. (**Io 5211**, 673 µm long); figs. 2, 3, ♀ RV, (holotype, **Io 5209**, 678 µm long); fig. 2, int. lat.; fig. 3, musc. sc. Scale A (250 µm; x 101), figs. 1, 2; scale B (100 µm; x 404), fig. 3.











ON *PARACYTHEREIDEA HEXALPHA* DORUK sp. nov.

by Neriman Doruk  
(Ege University, Izmir, Turkey)

*Paracytheridea hexalpha* sp. nov.

- 1971 *Paracytheridea* sp. 6, P. J. Barbeito-Gonzalez, *Mitt. Hamburg, zool. Mus. Inst.*, Bd. 67, 305, pl. 30, figs. 1a, 2a.  
1972 *Paracytheridea* sp. A, H. Uffenorde, *Göttinger Arb. Geol. Paläont.*, 13, 87, pl. 9, fig. 10, pl. 12, fig. 9.

*Holotype*: Brit. Mus. (Nat. Hist.) no. Io 5213, ♀ LV.

[Paratypes: Brit. Mus. (Nat. Hist.) nos. Io 5212, 5214, 5215].

*Type locality*: Drillings off south coast of Turkey; approx. lat. 36° 26' N, long. 35° 04' E. Pleistocene.*Derivation of name*: Greek, six + letter *alpha*, with reference to synonymy.

## Explanation of Plate 7, 148

Fig. 1, ♀ RV, ext. lat. (paratype, Io 5212, 890 µm long); fig. 2, ♀ LV, ext. lat. (holotype, Io 5213, 823 µm long); fig. 3, surface ornament (paratype, Io 5214).

Scale A (250 µm; x 90), fig. 1; scale B (250 µm; x 96), fig. 2; scale C (100 µm; x 282), fig. 3.

*Figured specimens*: Brit. Mus. (Nat. Hist.) nos. Io 5212 (♀ RV: Pl. 7, 148, fig. 1), Io 5213 (holotype, ♀ LV: Pl. 7, 148, fig. 2; Pl. 7, 150, fig. 1), Io 5214 (LV: Pl. 7, 148, fig. 3), Io 5215 (♂ RV: Pl. 7, 150, figs. 2, 3). Io 5212 and 5213 from the type locality, depth 450 - 480ft. — presumed shallow marine. Io 5214 and Io 5215 from Urla Bay (west coast of Turkey), about 30ft. depth. Subrecent, presumed littoral; approx. lat. 38° 19' N, long. 26° 47' E.

*Diagnosis*: Papillate solae with mural ghosts.



*Remarks*: Close to *Paracytheridea bovetensis* (Seguenza) but differs by having a weaker reticulation and more attenuated posterior processes. Sexual dimorphism is indicated by the presence of more elongate males.

*Distribution*: Recent of the Adriatic Sea and Pleistocene to Recent from Turkey.

## Explanation of Plate 7, 150

Fig. 1, ♀ LV, int. lat. (holotype, Io 5213, 823 µm long); fig. 2, ♂ RV, int. lat. (paratype, Io 5215, 723 µm long); fig. 3, ♂ RV, musc. sc. (paratype, Io 5215).

Scale A (250 µm; x 90), fig. 1; scale B (250 µm; x 96), fig. 2; scale C (100 µm; x 282), fig. 3.











ON *LEPTOCY THERE MULTIPUNCTATA* (SEGUENZA)

by Neriman Doruk  
(Ege University, Izmir, Turkey)

*Leptocythere multipunctata* (Seguenza, 1884)

- 1884 *Cythere multipunctata* G. Seguenza, *Il Naturalista Siciliana*, 29, pl. 1, figs. 9a - b.  
1950 *Leptocythere multipunctata* (Seguenza); G. Ruggieri, *Giorn. Geol. ser. 2*, 21, 52, pl. 1, figs. 8, 15.  
1972 *Leptocythere multipunctata multipunctata* (Seguenza); W. Sissingh, *Utrecht Micropaleont. Bull.*, 6, 91, pl. 5, fig. 9.

*Holotype*: Probably lost during the Messina earthquake of 1908 (Ruggieri, *Boll. Soc. Paleont.*, 2, no. 1, 3, 1963).

*Type locality*: Rizzola, near Syracuse, Sicily; Calabrian, Quaternary.

*Figured specimens*: Brit. Mus. (Nat. Hist.) nos. **Io 5180** (♀ RV: Pl. 7, 152, fig. 1; Pl. 7, 154, fig. 2), **Io 5181** (♂ LV: Pl. 7, 152, fig. 2; Pl. 7, 154, figs. 1, 3). Both are from drillings off Iskenderun Bay, S Turkey at a depth of 610 feet below sea floor, approx. lat. 36° 07' N, long. 36° 13' E; Pliocene - Pleistocene; presumed shallow marine.

*Diagnosis*: Carapace with diagnostic outline; shell surface irregularly pitted with numerous puncta.

Explanation of Plate 7, 152


Fig. 1, ♀ RV, ext. lat. (**Io 5180**, 470 µm long); fig. 2, ♂ LV, ext. lat. (**Io 5181**, 450 µm long).  
Scale A (250 µm; x 173), fig. 1; scale B (250 µm; x 188), fig. 2.

*Remarks*: Some authors have distinguished 2 subspecies of *L. multipunctata* on the size of the surface puncta (e.g. Sissingh 1972, *op. cit.*) but as a certain amount of variation in puncta size is seen in the present material from the same sample, these are here considered to be one species.

Sexual dimorphism is only slightly observed in this species with females being slightly higher than the males. Marginal pore canals branching (see text-fig. 1).

*Distribution*: Recent: Adriatic, Porto-Corsini (Ruggieri, 1950, *op. cit.*). Quaternary: Syracuse, Sicily (Seguenza, 1884, *op. cit.*); Imola, Italy (Ruggieri, 1950). Upper Pliocene - Pleistocene: Aegean Islands (Sissingh, 1972, *op. cit.*) and S Turkey (present paper).



Text-fig. 1. Marginal pore canals  
of *L. multipunctata*, ♀ RV, **Io 5180** (x 66).



Explanation of Plate 7, 154

Figs. 1, 3, ♂ LV (**Io 5181**, 450 µm long): fig. 1, int. lat.; fig. 3, musc. sc.; fig. 2, ♀ RV, int. lat. (**Io 5180**, 470 µm long).  
Scale A (250 µm; x 159), fig. 1; scale B (250 µm; x 148), fig. 2; scale C (100 µm; x 477), fig. 3.









## General Index

- Acanthocythereis decoris* Siddiqui; 131 - 134  
 Athersuch, J., On *Aurila woodwardii* (Brady); 45 - 52  
 Athersuch, J., On *Loxoconcha pontica* Klie; 53 - 60  
 Athersuch, J. & Whittaker, J. E., On *Callistocythere crispata* (Brady); 67 - 72  
 Athersuch, J. & Whittaker, J. E., On *Callistocythere littoralis* (Müller); 61 - 66  
*audax*, *Scottia*; 37 - 44  
*Aurila woodwardii* (Brady); 45 - 52  
 Bate, R. H. & Sheppard, L. M., On *Quadraleberis exquisita* Bate & Sheppard gen. et sp. nov.; 107 - 112  
 Bate, R. H. & Sheppard, L. M., On *Xestoleberis postangulata* Bate & Sheppard sp. nov.; 101 - 106  
*bilobata*, *Schuleridea*; 25 - 28  
*braderupensis*, *Pachydomelloides*; 77 - 80  
*Bulbosclerites unicornis* (Neckaja); 73 - 76  
*Callistocythere crispata* (Brady); 67 - 72  
*Callistocythere littoralis* (Müller); 61 - 66  
*Callistocythere mediterranea* (Müller); 135 - 138  
*Callistocythere montana* Doruk sp. nov.; 139 - 142  
*crispata*, *Callistocythere*; 67 - 72  
*Cyprideis exuberans* van Harten sp. nov.; 89 - 100  
*Cyprideis undosa* van Harten sp. nov.; 81 - 88  
 DeDeckker, P., On *Scottia audax* (Chapman); 37 - 44  
*decoris*, *Acanthocythereis*; 131 - 134  
*Disulcina syltensis* (Schallreuter); 21 - 24  
 Doruk, N., On *Callistocythere mediterranea* (Müller); 135 - 138  
 Doruk, N., On *Callistocythere montana* Doruk sp. nov.; 139 - 142  
 Doruk, N., On *Leptocythere multipunctata* (Seguenza); 151 - 154  
 Doruk, N., On *Paracytheridea hexalpha* Doruk sp. nov.; 147 - 150  
 Doruk, N., On *Paracytheridea inscita* Doruk sp. nov.; 143 - 146  
*Eographiodactylus sulcatus* Schallreuter; 1 - 8  
*exquisita*, *Quadraleberis*; 107 - 112  
*exuberans*, *Cyprideis*, 89 - 100  
*Foramenella parkis* (Neckaja); 17 - 20  
 Fuller, N., On *Procytheropteron prolongatum* (Sharapova); 117 - 124  
 Harten, D. van, On *Cyprideis exuberans* van Harten sp. nov.; 89 - 100  
 Harten, D. van, On *Cyprideis undosa* van Harten sp. nov.; 81 - 88  
*hexalpha*, *Paracytheridea*; 147 - 150  
 Horne, D. J. & Whittaker, J. E., On *Semicytherura tela* Horne & Whittaker sp. nov.; 125 - 130  
*inscita*, *Paracytheridea*; 143 - 146  
*Klimphores planus* Schallreuter; 9 - 16  
*Leptocythere multipunctata* (Seguenza); 151 - 154  
*littoralis*, *Callistocythere*; 61 - 66  
*Loxoconcha pontica* Klie; 53 - 60  
*mediterranea*, *Callistocythere*; 135 - 138  
*Micropneumatocythere tumida* Stephens & Ware sp. nov.; 113 - 116  
*montana*, *Callistocythere*; 139 - 142  
*multipunctata*, *Leptocythere*; 151 - 154  
 Neale, J. W., On *Paranotacythere speetonensis* (Neale); 33 - 36  
 Neale, J. W., On *Schuleridea bilobata* (Triebe); 25 - 28  
 Neale, J. W., On *Schuleridea rhomboidalis* Neale; 29 - 32  
*Pachydomelloides braderupensis* Schallreuter sp. nov.; 77 - 80  
*Paracytheridea hexalpha* Doruk sp. nov.; 147 - 150  
*Paracytheridea inscita* Doruk sp. nov.; 143 - 146  
*Paranotacythere speetonensis* (Neale); 33 - 36

- parkis*, *Foramenella*; 17 - 20  
*planus*, *Klimphores*; 9 - 16  
*pontica*, *Loxoconcha*; 53 - 60  
*postangulata*, *Xestoleberis*; 101 - 106  
*Procytheropteron prolongatum* (Sharapova); 117 - 124  
*prolongatum*, *Procytheropteron*; 117 - 124  
*Quadraleberis exquisita* Bate & Sheppard gen. et sp. nov.; 107 - 112  
*rhomboidalis*, *Schuleridea*; 29 - 32  
Schallreuter, R. E. L., On *Bulbosclerites unicornis* (Neckaja); 73 - 76  
Schallreuter, R. E. L., On *Disulcina syltensis* Schallreuter sp. nov.; 21 - 24  
Schallreuter, R. E. L., On *Eographiodactylus sulcatus* Schallreuter; 1 - 8  
Schallreuter, R. E. L., On *Foramenella parkis* (Neckaja); 17 - 20  
Schallreuter, R. E. L., On *Klimphores planus* Schallreuter; 9 - 16  
Schallreuter, R. E. L., On *Pachydomelloides braderupensis* Schallreuter sp. nov.; 77 - 80  
*Schuleridea bilobata* (Triebel); 25 - 28  
*Schuleridea rhomboidalis* Neale; 29 - 32  
*Scottia audax* (Chapman); 37 - 44  
*Semicytherura tela* Horne & Whittaker sp. nov.; 125 - 130  
Sheppard, L. M. & Bate, R. H., On *Quadraleberis exquisita* Bate & Sheppard gen. et sp. nov.; 107 - 112  
Sheppard, L. M. & Bate, R. H., On *Xestoleberis postangulata* Bate & Sheppard sp. nov.; 101 - 106  
Siddiqui, Q. A., On *Acanthocythereis decoris* Siddiqui; 131 - 134  
*speetonensis*, *Paranotacythere*; 33 - 36  
Stephens, J. & Ware, N., On *Micropneumatocythere tumida* Stephens & Ware sp. nov.; 113 - 116  
*sulcatus*, *Eographiodactylus*; 1 - 8  
*syltensis*, *Disulcina*; 21 - 24  
*tela*, *Semicytherura*; 125 - 130  
*tumida*, *Micropneumatocythere*; 113 - 116  
*undosa*, *Cyprideis*; 81 - 88  
*unicornis*, *Bulbosclerites*; 73 - 76  
Ware, M. & Stephens, J., On *Micropneumatocythere tumida* Stephens & Ware sp. nov.; 113 - 116  
Whittaker, J. E. & Athersuch, J., On *Callistocythere crispata* (Brady); 67 - 72  
Whittaker, J. E. & Athersuch, J., On *Callistocythere littoralis* (Müller); 61 - 66  
Whittaker, J. E. & Horne, D. J., On *Semicytherura tela* Horne & Whittaker sp. nov.; 125 - 130  
*woodwardii*, *Aurila*; 45 - 52  
*Xestoleberis postangulata* Bate & Sheppard sp. nov.; 101 - 106

## Index; Geological Horizon

See 1 (2) 5 - 22 (1973) for explanation of the Schedules in the Universal Decimal Classification

- |   |   |
|---|---|
| (113.31) Ordovician:                            | (116.311) Berriasian:                         |
| <i>Foramenella parkis</i> ; 17 - 22             | <i>Paranotacythere speetonensis</i> ; 33 - 36 |
| (113.312) Middle Ordovician:                    | (116.312) Aptian:                             |
| <i>Klimphores planus</i> ; 9 - 16               | <i>Schuleridea bilobata</i> ; 25 - 28         |
| (113.313) Upper Ordovician:                     | <i>Schuleridea rhomboidalis</i> ; 29 - 32     |
| <i>Bulbosclerites unicornis</i> ; 73 - 76       | (118.14) Eocene:                              |
| <i>Disulcina syltensis</i> ; 21 - 24            | <i>Acanthocythereis decoris</i> ; 131 - 134   |
| <i>Eographiodactylus sulcatus</i> ; 1 - 8       | (118.21) Miocene:                             |
| (116.222) Bathonian:                            | <i>Cyprideis exuberans</i> ; 89 - 100         |
| <i>Micropneumatocythere tumida</i> ; 113 - 116  | <i>Cyprideis undosa</i> ; 81 - 88             |
| (116.223) Volgian;                              | (118.213) Tortonian:                          |
| <i>Procytheropteron prolongatum</i> ; 117 - 124 | <i>Callistocythere montana</i> ; 139 - 142    |
|   | <i>Paracytheridea inscisa</i> ; 143 - 146     |



- |   |  |
|---|--|
| <p>(118.22) Pliocene:<br/> <i>Callistocythere mediterranea</i>; 135 - 138<br/> <i>Cyprideis exuberans</i>; 89 - 100<br/> <i>Leptocythere multipunctata</i>; 151 - 154</p> <p>(119) Quaternary:<br/> <i>Aurila woodwardii</i>; 45 - 52<br/> <i>Loxoconcha pontica</i>; 53 - 60</p> <p>(119.1) Pleistocene:<br/> <i>Leptocythere multipunctata</i>; 151 - 154<br/> <i>Paracytheridea hexalpa</i>; 147 - 150</p> | <p>(119.4) Holocene:<br/> <i>Paracytheridea hexalpa</i>; 147 - 150</p> <p>(119.9) Recent:<br/> <i>Callistocythere crispata</i>; 67 - 72<br/> <i>Callistocythere littoralis</i>; 61 - 66<br/> <i>Callistocythere mediterranea</i>; 135 - 138<br/> <i>Leptocythere multipunctata</i>; 151 - 154<br/> <i>Paracytheridea hexalpa</i>; 147 - 150<br/> <i>Quadraleberis exquisita</i>; 107 - 112<br/> <i>Scottia audax</i>; 37 - 44<br/> <i>Semicytherura tela</i>; 125 - 130<br/> <i>Xestoleberis postangulata</i>; 101 - 106</p> |
|---|--|

### Index; Geographical Location

See 1 (2) 5 - 22 (1973) for explanation of the Schedules in the Universal Decimal Classification

- |  |  |
|--|--|
| <p>(261.2) NE Atlantic:<br/> <i>Callistocythere crispata</i>; 67 - 72<br/> <i>Callistocythere littoralis</i>; 61 - 66<br/> <i>Semicytherura tela</i>; 125 - 130</p> <p>(262) Mediterranean Sea:<br/> <i>Aurila woodwardii</i>; 45 - 52<br/> <i>Loxoconcha pontica</i>; 53 - 60</p> <p>(267.8) Persian Gulf:<br/> <i>Quadraleberis exquisita</i>; 107 - 112<br/> <i>Xestoleberis postangulata</i>; 101 - 106</p> <p>(420) England:<br/> <i>Micropneumatocythere tumida</i>; 113 - 116</p> <p>(427.4) Yorkshire:<br/> <i>Paranotacythere speetonensis</i>; 33 - 36<br/> <i>Schuleridea bilobata</i>; 25 - 28<br/> <i>Schuleridea rhomboidalis</i>; 29 - 32</p> <p>(430.1) Federal Republic of Germany:<br/> <i>Schuleridea bilobata</i>; 25 - 28<br/> <i>Schuleridea rhomboidalis</i>; 29 - 32</p> <p>(430.2) German Democratic Republic:<br/> <i>Klimphores planus</i>; 9 - 16</p> <p>(45) Italy:<br/> <i>Callistocythere mediterranea</i>; 135 - 138</p> | <p>(46) Spain:<br/> <i>Cyprideis exuberans</i>; 89 - 100<br/> <i>Cyprideis undosa</i>; 81 - 88</p> <p>(47) U.S.S.R.:<br/> <i>Procytheropteron prolongatum</i>; 117 - 124</p> <p>(486) Gotland:<br/> <i>Eoglyphiodactylus sulcatus</i>; 1 - 8<br/> <i>Foramenella parkis</i>; 17 - 20</p> <p>(489) Denmark:<br/> <i>Schuleridea bilobata</i>; 25 - 28</p> <p>(492) Netherlands:<br/> <i>Bulbosclerites unicornis</i>; 73 - 76<br/> <i>Disulcina syltensis</i>; 21 - 24<br/> <i>Eoglyphiodactylus sulcatus</i>; 1 - 8<br/> <i>Foramenella parkis</i>; 17 - 29<br/> <i>Pachydomelloides braderupensis</i>; 77 - 80</p> <p>(549) Pakistan:<br/> <i>Acanthocythereis decoris</i>; 113 - 116</p> <p>(560) Turkey (Asia):<br/> <i>Callistocythere mediterranea</i>; 135 - 138<br/> <i>Callistocythere montana</i>; 139 - 142<br/> <i>Leptocythere multipunctata</i>; 151 - 154<br/> <i>Paracytheridea hexalpa</i>; 147 - 150<br/> <i>Paracytheridea inscita</i>; 143 - 146</p> <p>(931) New Zealand:<br/> <i>Scottia audax</i>; 37 - 44</p> |
|--|--|



# A Stereo-Atlas of Ostracod Shells

edited by R. H. Bate, J. W. Neale, Lesley M. Sheppard  
and David J. Siveter

Volume 7, 1980

Part 1 (pp.1–72); 27th June, 1980

Part 2 (pp.73–157); 19th December, 1980

Published by The British Micropalaeontological Society  
in association with Robertson Research International Ltd., Llandudno, Wales



## Contents

1	On <i>Eographiodactylus sulcatus</i> Schallreuter; by R. E. L. Schallreuter	1
2	On <i>Klimphores planus</i> Schallreuter; by R. E. L. Schallreuter	9
3	On <i>Foramenella parkis</i> (Neckaja); by R. E. L. Schallreuter	17
4	On <i>Disulcina syltensis</i> Schallreuter sp. nov.; by R. E. L. Schallreuter	21
5	On <i>Schuleridea bilobata</i> (Triebel); by J. W. Neale	25
6	On <i>Schuleridea rhomboidalis</i> Neale; by J. W. Neale	29
7	On <i>Paranotacythere speetonensis</i> (Neale); by J. W. Neale	33
8	On <i>Scottia audax</i> (Chapman); by P. De Deckker	37
9	On <i>Aurila woodwardii</i> (Brady); by J. Athersuch	45
10	On <i>Loxoconcha pontica</i> Klie; by J. Athersuch	53
11	On <i>Callistocythere littoralis</i> (Müller); by J. Athersuch & J. E. Whittaker	61
12	On <i>Callistocythere crispata</i> (Brady); by J. Athersuch & J. E. Whittaker	67
13	On <i>Bulbosclerites unicornis</i> (Neckaja); by R. E. L. Schallreuter	73
14	On <i>Pachydomelloides braderupensis</i> Schallreuter sp. nov.; by R. E. L. Schallreuter	77
15	On <i>Cyprideis undosa</i> van Harten sp. nov.; by D. van Harten	81
16	On <i>Cyprideis exuberans</i> van Harten sp. nov.; by D. van Harten	89
17	On <i>Xestoleberis postangulata</i> Bate & Sheppard sp. nov.; by R. H. Bate & L. M. Sheppard	101
18	On <i>Quadraleberis exquisita</i> Bate & Sheppard gen. et sp. nov.; by R. H. Bate & L. M. Sheppard	107
19	On <i>Micropneumatocythere tumida</i> Stephens & Ware sp. nov.; by J. Stephens & M. Ware	113
20	On <i>Procytheropteron prolongatum</i> (Sharapova); by N. Fuller	117
21	On <i>Semicytherura tela</i> Horne & Whittaker sp. nov.; by D. J. Horne & J. E. Whittaker	125
22	On <i>Acanthocythereis decoris</i> Siddiqui; by Q. A. Siddiqui	131
23	On <i>Callistocythere mediterranea</i> (Müller); by N. Doruk	135
24	On <i>Callistocythere montana</i> Doruk sp. nov.; by N. Doruk	139
25	On <i>Paracytheridea inscita</i> Doruk sp. nov.; by N. Doruk	143
26	On <i>Paracytheridea hexalpha</i> Doruk sp. nov.; by N. Doruk	147
27	On <i>Leptocythere multipunctata</i> (Seguenza); by N. Doruk	151
28	Index for Volume 7, 1980	155

# A stratigraphical index of British Ostracoda

*The first of a series of Occasional Papers of the British Micropalaeontological Society. ISBN 0 902354 08 6*

*Edited by Raymond H. Bate and Eric Robinson*

**Geological Journal      Special Issue No. 8**

Developed from ostracod specialist meetings of the British Micropalaeontological Society this book aims to illustrate the stratigraphically important index ostracods of Britain. Brief descriptions of the ostracods, together with palaeogeographic maps, range tables, discussions on present knowledge and on faunal associations make this a valuable book for both student and professional micropalaeontologist. Introductory chapters on the biography of past ostracod workers and on ostracod morphology add to the book's value as a reference work. 500 pp

## Contents

Editorial Preface	
History of ostracod studies	<i>Eric Robinson</i>
Ostracod morphology	<i>Raymond H. Bate and Eric Robinson</i>
The Ordovician	<i>David Siveter</i>
The Silurian	<i>David Siveter</i>
The Devonian	<i>Andrew Gooday</i>
The Carboniferous	<i>Eric Robinson</i>
The Permian	<i>Eric Robinson</i>
The Trias	<i>Raymond H. Bate</i>
The Jurassic Pt I — Pliensbachian to Toarcian	<i>Alan Lord</i>
The Jurassic Pt II — Aalenian to Bathonian	<i>Raymond H. Bate</i>
The Jurassic Pt III — Callovian to Portlandian	<i>Tom Kilenyi</i>
The Purbeck/Wealden	<i>Tom Kilenyi and John W. Neale</i>
The Cretaceous	<i>John W. Neale</i>
The Tertiary — Palaeogene	<i>Mike Keen</i>
The Pleistocene	<i>Eric Robinson</i>
Ostracod range tables — commercial application	<i>Lesley M. Sheppard</i>
Index	

*Published 1978*

*Price, including postage, £40.00 (U.S. \$112.90)*

Available to members of the British Micropalaeontological Society at the listed price less 25%.

*(Attn. John Wilson), John Wiley and Sons Ltd., Baffins Lane, Chichester, Sussex PO19 1UD, England.*

# Palaeontological microslides



## EK Hull Microslide Company

24 Lynmouth Gardens  
Perivale, Middlesex UB6 7HR  
England  
Telephone 01 - 998 2256

### Cardboard slides in aluminium holders

Glass or thick, clear,  
acetate coverslides.  
Single, 4-celled or faunal  
(32 or 64 cell divisions)

### Plastic slides

Single, double, 3 or 4-celled  
and faunal  
(32 or 64 cell divisions)  
All slides sold complete  
at 20 pence each  
(USA 47 cents)

For air-freight, sold  
without glass coverslides  
at 19 pence  
(USA 45 cents)

Postage and packing extra

# Stereo - Atlas of Ostracod Shells: Vol.7, Part 2

## CONTENTS

7 (13) 73 - 76	On <i>Bulbosclerites unicornis</i> (Neckaja); by R. E. L. Schallreuter	(£1.40)
7 (14) 77 - 80	On <i>Pachydomelloides braderupensis</i> Schallreuter sp. nov.; by R. E. L. Schallreuter	(£1.40)
7 (15) 81 - 88	On <i>Cyprideis undosa</i> van Harten sp. nov.; by D. van Harten	(£2.80)
7 (16) 89 - 100	On <i>Cyprideis exuberans</i> van Harten sp. nov.; by D. van Harten	(£4.20)
7 (17) 101 - 106	On <i>Xestoleberis postangulata</i> Bate & Sheppard sp. nov.; by R. H. Bate & L. M. Sheppard	(£2.10)
7 (18) 107 - 112	On <i>Quadraleberis exquisita</i> Bate & Sheppard gen. et sp. nov.; by R. H. Bate & L. M. Sheppard	(£2.10)
7 (19) 113 - 116	On <i>Micropneumatocythere tumida</i> Stephens & Ware sp. nov.; by J. Stephens & M. Ware	(£1.40)
7 (20) 117 - 124	On <i>Procytheropteron prolongatum</i> (Sharapova); by N. Fuller	(£2.80)
7 (21) 125 - 130	On <i>Semicytherura tela</i> Horne & Whittaker sp. nov.; by D. J. Horne & J. E. Whittaker	(£2.10)
7 (22) 131 - 134	On <i>Acanthocythereis decoris</i> Siddiqui; by Q. A. Siddiqui	(£1.40)
7 (23) 135 - 138	On <i>Callistocythere mediterranea</i> (Müller); by N. Doruk	(£1.40)
7 (24) 139 - 142	On <i>Callistocythere montana</i> Doruk sp. nov.; by N. Doruk	(£1.40)
7 (25) 143 - 146	On <i>Paracytheridea inscita</i> Doruk sp. nov.; by N. Doruk	(£1.40)
7 (26) 147 - 150	On <i>Paracytheridea hexalpha</i> Doruk sp. nov.; by N. Doruk	(£1.40)
7 (27) 151 - 154	On <i>Leptocythere multipunctata</i> (Seguenza); by N. Doruk	(£1.40)
7 (28) 155 - 157	Index for Volume 7, 1980	(£0.60)

Prepaid annual subscription (valid for Volume 8, 1981)

Individual subscription £20.00 or US \$ 50.00 for 2 parts (post free)

Price per Part: £20.00 or US \$ 50.00

Institutional subscription £30.00 or US \$ 75.00 for 2 parts (post free)

Price per Part: £30.00 or US \$ 75.00

Back volumes: Vol. 1 (4 Parts): £20.00; price per Part: £5.00

Vol. 2 (4 Parts): £28.00; price per Part: £7.00

Vol. 3 (2 Parts): £24.00; price per Part: £12.00

Vol. 4 (2 Parts): £30.00; price per Part: £15.00

Vol. 5 (2 Parts): £32.00; price per Part: £16.00

Vol. 6 (2 Parts): £40.00; price per Part: £20.00

Vol. 7 (2 Parts): £40.00; price per Part: £20.00

Prices of Numbers as shown in Lists of Contents for each Part

Postage extra in sales of all back Numbers and back Parts

No trade discount is allowed on the subscription rate

Orders should be addressed to: Dr. C. W. Haskins,

Robertson Research International Limited,

'Ty'n-y-Coed', Llanrhos, Llandudno, Gwynedd LL30 1SA.

Cheques should be made payable to Robertson Research Co. (Stereo-Atlas Account)

## SPECIAL OFFER

*50% off all back part prices if*

*you become a subscriber to the Atlas*